arXiv:1601.04320v2 [math.QA] 5 Feb 2016

DOUBLE-BOSONIZATION AND MAJID'S CONJECTURE, (III): TYPE-CROSSING AND INDUCTIONS OF E_6 AND E_7, E_8

HONGMEI HU AND NAIHONG HU*

Abstract. Double-bosonization construction in Majid [M1] is expectedly allowed to generate a tree of quantum groups. Some main branches of the tree in [HH1, HH2] have been depicted how to grow up. This paper continues to elucidate the type-crossing and inductive constructions of exceptional quantum groups of types E_6 and E_7, E_8, respectively, based on the generalized double-bosonization Theorem established in [HH2]. Thus the Majid's expectation for the inductive constructions of $U_q(\mathfrak{g})$'s for all finite-dimensional complex simple Lie algebras is completely achieved.

1. Introduction and our results

A striking feature of quantum group theory is in close connection with many branches of mathematics and physics, such as Lie groups, Lie algebras and their representations, Hecke algebras and their representation theory, quantum invariants theory of knots or links and 3-manifolds, as well as the current hot studies on monoidal categories and various categorifications, etc. So quantum group theory always attracts many mathematicians to find some better ways in a suitable framework to construct their structures defined initially by generators and relations. In the early nineties, Ringel [Ri] realized the positive part of quantum groups by quiver representations and Hall algebras, which inspired Lusztig's canonical base theory [L1, L2]. Bridgeland [Br] realized the entire quantum groups of type *ADE* by the Ringel-Hall algebras. In [Ro], Rosso also realized the positive part of $U_q(\mathfrak{g})$ by the quantum shuffle in a braided category, and gave a recipe on inductive constructions in the sense of quantum shuffle. The axiomatic construction for the entire quantum groups for the standard Drinfeld-Jimbo types (cf. Fang-Rosso in [FR]) has been nontrivially generalized to the multi-parameter setting by Hu-Li-Rosso [HLR]. On the other hand, Majid [M1] (also Sommerhäuser in [So]) constructed $U_q(\mathfrak{g})$ by establishing the double-bosonization theory in a braided category of H-modules (resp. in a larger braided category, i.e., the category of Yetter-Drinfeld H-modules). In this paper, we mainly focus on the double-bosonization method in [M1].

*N. H., supported by the NSFC (Grant No. 11271131).

Associated with any mutually dual braided groups $B^\star, B$ covariant under a background quasitriangular Hopf algebra H, there is a new quantum group structure on the tensor space $B^\star \otimes H \otimes B$ by double-bosonization in [M1], consisting of H extended by B as additional 'positive roots' and its dual $B^\star$ as additional 'negative roots'. The construction is more direct than the quantum double because one can reach to $U_q(\mathfrak{g})$ (rather than to take a quotient from the quantum double). Specially, Majid regarded $U_q(\mathfrak{n}^\pm)$ as the mutually dual braided groups in the braided category of left H-modules for 'Cartan subalgebra' H, then recovered $U_q(\mathfrak{g})$ by double-bosonization. On the other hand, Majid claimed that the double-bosonization construction allowed to be used to generate a tree of quantum groups. That is, many new quantum groups and the inductive constructions of $U_q(\mathfrak{g})$'s for all complex simple Lie algebras $\mathfrak{g}$ were supposed to be obtained by this theory in [M1]. He thought that at each node of the tree, there are many choices to adjoin a pair of certain braided groups covariant under the quantum group at that node. This is a representation-theoretical challenge to elaborate the full tree structure. After a few examples given by Majid himself [M1, M6] twenty years ago, recently, almost main branches have been depicted in [HH1, HH2, HH3]. The remaining situation left is the E-series. This consists of the aim of this paper.

Nowadays, Majid's framework on braided groups in certain braided categories has been developed into the framework on Nichols algebras in the Yetter-Drinfeld categories for the purpose to classify finite-dimensional pointed Hopf algebras; see the works [AS1, AS2] of Andruskiewitsch and Schneider, and the one [HS] of Heckenberger and Schneider. Hence, we believe in some sense that it is significant to study what kinds of new finite-dimensional Hopf algebras can be found via the double-bosonization procedure. An interesting application of our constructions might be connected to a recent work of Cuntz and Lentner on Nichols algebras (also see the concluding remarks in [G] for further interesting considerations).

The paper is organized as follows. In section 2, we recall some basic facts about FRT-bialgebras associated with R-matrices in [FRT], Majid's double bosonization theorem in [M1] and the generalized double-bosonization construction theorem established in [HH2] suitable for those irregular R-matrices we encountered when we treat the exceptional cases (beyond the considerations in [M1]). In view of some results and methods in [HH1, HH2], we construct $U_q(E_6)$ by working on $U_q(D_5)$ and its one of the half-spin representations, $U_q(E_7)$, $U_q(E_8)$ can be constructed from $U_q(E_6)$, $U_q(E_7)$ and their minimal dimensional fundamental representations in section 3. These representations are given in diagrams (see Figures 1 to 3). In the last section, we give a proposition to summarize our inductive constructions of $U_q(\mathfrak{g})$'s for all complex finite-dimensional simple Lie algebras, and also reflect

these constructions on the Dynkin diagrams. With these results, we see that both the double-bosonization theory and the quantum shuffle theory (due to Rosso [Ro]) yield the same tree structure of $U_q(\mathfrak{g})$'s for all complex semisimple Lie algebras.

2. Preliminaries

In this paper, let k be complex field, $\mathbb{R}$ real field, E the Euclidean space $\mathbb{R}^n$ or its a suitable subspace. ε_i denotes the usual orthogonal unit vectors in $\mathbb{R}^n$. $\mathfrak{g}$ is a finite-dimensional complex semisimple Lie algebra with simple roots α_i. λ_i is the fundamental weight corresponding to simple root α_i. Cartan matrix of $\mathfrak{g}$ is (a_{ij}), where $a_{ij} = \frac{2(\alpha_i,\alpha_j)}{(\alpha_i,\alpha_i)}$, and $d_i = \frac{(\alpha_i,\alpha_i)}{2}$. Let $(H, \mathscr{R})$ be a quasitriangular Hopf algebra, where $\mathscr{R}$ is called the universal R-matrix, $\mathscr{R} = \mathscr{R}^{(1)} \otimes \mathscr{R}^{(2)}$, $\mathscr{R}_{21} = \mathscr{R}^{(2)} \otimes \mathscr{R}^{(1)}$, we denote by Δ, η, ϵ its coproduct, counit, unit, and by S its antipode.

We shall use Sweedler's notation: for $h \in H$, $\Delta(h) = h_1 \otimes h_2$. $H^{\rm op}$ ($H^{\rm cop}$) denotes the opposite (co)algebra structure of H, respectively. $\mathfrak{M}_H$ (${}_H\mathfrak{M}$) denotes the braided category consisting of right (left) H-modules, respectively. If there is a coquasitriangular Hopf algebra A such that dual pair (H, A) is a weakly quasitriangular, then $\mathfrak{M}_H$ (${}_H\mathfrak{M}$) is equivalent to the braided category ${}^A\mathfrak{M}$ ($\mathfrak{M}^A$) of left (right) A-comodules, respectively. For the detailed description of these theories, we left to the readers to refer to Drinfeld's and Majid's papers [D], [M2], [M3], and so on. Braided group is a braided bialgebra or Hopf algebra in some braided category, in order to distinguish from the ordinary Hopf algebras, let $\underline{\Delta}$, $\underline{S}$ denote its coproduct and antipode, respectively.

2.1. **Majid's double-bosonization theory.** Let C, B be a pair of braided groups in $\mathfrak{M}_H$, which are called dually paired if there is an intertwiner ev : $C \otimes B \longrightarrow k$ such that

$$\mathrm{ev}(cd, b) = \mathrm{ev}(d, \underline{b_{(1)}})\mathrm{ev}(c, \underline{b_{(2)}}), \quad \mathrm{ev}(c, ab) = \mathrm{ev}(\underline{c_{(2)}}, a)\mathrm{ev}(\underline{c_{(1)}}, b), \quad \forall a, b \in B, c, d \in C.$$

Then $C^{\rm op/cop}$(with opposite product and coproduct) is a Hopf algebra in ${}_H\mathfrak{M}$, which is dual to B in the sense of an ordinary duality pairing $\langle\ ,\ \rangle$, which is H-bicovariant: $\langle h \triangleright c, b\rangle = \langle c, b \triangleleft h\rangle$, for all $h \in H$. Let $\overline{C} = (C^{\rm op/cop})^{\underline{\rm cop}}$, then $\overline{C}$ is a braided group in ${}_{\overline{H}}\overline{\mathfrak{M}}$, where $\overline{H}$ is $(H, \mathscr{R}_{21}^{-1})$. With these, Majid gave the following double bosonization theorem and some results in [M1]:

Theorem 2.1. (**Majid**) *On the tensor space $\overline{C} \otimes H \otimes B$, there is a unique Hopf algebra structure $U = U(\overline{C}, H, B)$ such that $H \ltimes B$ and $\overline{C} \rtimes H$ are sub-Hopf algebras by the canonical inclusions and*

$$bc = (\mathscr{R}_1^{(2)} \triangleright c_{\overline{(2)}})\mathscr{R}_2^{(2)}\mathscr{R}_1^{-(1)}(b_{\underline{(2)}} \triangleleft \mathscr{R}_2^{-(1)})\,\langle \mathscr{R}_1^{(1)} \triangleright c_{\overline{(1)}}, b_{\underline{(1)}} \triangleleft \mathscr{R}_2^{(1)}\rangle \cdot$$
$$\cdot\,\langle \mathscr{R}_1^{-(2)} \triangleright \overline{S}\, c_{\overline{(3)}}, b_{\underline{(3)}} \triangleleft \mathscr{R}_2^{-(2)}\rangle,$$

for all $b \in B$, $c \in \overline{C}$ viewed in U. Here $\mathscr{R}_1, \mathscr{R}_2$ are distinct copies of the quasitriangular structure $\mathscr{R}$ of H. The product, coproduct of U are given by

$$(c \otimes h \otimes b) \cdot (d \otimes g \otimes a) = c(h_{(1)}\mathscr{R}_1^{(2)} \triangleright d_{\overline{(2)}}) \otimes h_{(2)}\mathscr{R}_2^{(2)}\mathscr{R}_1^{-(1)} g_{(1)} \otimes (b_{\underline{(2)}} \triangleleft \mathscr{R}_2^{-(1)} g_{(2)})a$$
$$\langle \mathscr{R}_1^{(1)} \triangleright d_{\overline{(1)}}, b_{\underline{(1)}} \triangleleft \mathscr{R}_2^{(1)} \rangle \langle \mathscr{R}_1^{-(2)} \triangleright \overline{S} d_{\overline{(3)}}, b_{\underline{(3)}} \triangleleft \mathscr{R}_2^{-(2)} \rangle;$$

$$\Delta(c \otimes h \otimes b) = c_{\overline{(1)}} \otimes \mathscr{R}^{-(1)} h_{(1)} \otimes b_{\underline{(1)}} \triangleleft \mathscr{R}^{(1)} \otimes \mathscr{R}^{-(2)} \triangleright c_{\overline{(2)}} \otimes h_{(2)} \mathscr{R}^{(2)} \otimes b_{\underline{(2)}}.$$

Remark 2.1. *If there exists a coquasitriangular Hopf algebra A such that (H, A) is a weakly quasitriangular dual pair, and b, c are primitive elements, then some relations simplify to*

$$[b, c] = \mathscr{R}(b^{\overline{(1)}}) \langle c, b^{\overline{(2)}} \rangle - \langle c^{\overline{(1)}}, b \rangle \bar{\mathscr{R}}(c^{\overline{(2)}});$$

$$\Delta b = b^{\overline{(2)}} \otimes \mathscr{R}(b^{\overline{(1)}}) + 1 \otimes b, \quad \Delta c = c \otimes 1 + \bar{\mathscr{R}}(c^{\overline{(2)}}) \otimes c^{\overline{(1)}}.$$

2.2. **Generalized double-bosonization construction theorem.** Let R be an invertible matrix obeying the quantum Yang-Baxter equation. There is a bialgebra $A(R)$ [FRT] corresponding to the R-matrix, called an FRT-bialgebra.

Definition 2.1. *$A(R)$ is generated by 1 and $t = \{t^i_j\}$, having the following structure:*

$$R^{i\ j}_{a\ b} t^b_k t^a_l = t^i_a t^j_b R^{a\ b}_{l\ k}, \qquad \Delta(t^i_j) = t^i_a \otimes t^a_j, \quad \epsilon(t^i_j) = \delta^i_j.$$

$A(R)$ is a coquasitriangular bialgebra with $\mathscr{R} : A(R) \otimes A(R) \longrightarrow k$ such that $\mathscr{R}(t^i_j \otimes t^k_l) = R^{ik}_{jl}$. Here $T_1 = T \otimes I$, $T_2 = I \otimes T$, R^{ik}_{jl} denotes the entry at row (ik) and column (jl) in R.

By double cross product of bialgebra $A(R)$, Majid obtained bialgebra $\widetilde{U(R)}$ in [M5], which is generated by $m^{\pm}$ and satisfies

$$Rm^{\pm}_1 m^{\pm}_2 = m^{\pm}_2 m^{\pm}_1 R, \quad Rm^{+}_1 m^{-}_2 = m^{-}_2 m^{+}_1 R,$$
$$\Delta((m^{\pm})^i_j) = (m^{\pm})^a_j \otimes (m^{\pm})^i_a, \quad \epsilon((m^{\pm})^i_j) = \delta_{ij},$$

where $(m^{\pm})^i_j$ denotes the entry at row i and column j in $m^{\pm}$. Moreover, $(\widetilde{U(R)}, A(R))$ is a weakly quasitriangular dual pair of bialgebras. In braided category ${}^{A(R)}\mathfrak{M}$ ($\mathfrak{M}^{A(R)}$), there are two classical braided groups $V(R', R)$, $V^{\vee}(R', R^{-1}_{21})$ in [M4], called braided (co)vector algebras, respectively.

Proposition 2.1. *Suppose that R' is another matrix such that (i) $R_{12}R_{13}R'_{23} = R'_{23}R_{13}R_{12}$, $R_{23}R_{13}R'_{12} = R'_{12}R_{13}R_{23}$, (ii) $(PR + 1)(PR' - 1) = 0$, (iii) $R_{21}R'_{12} = R'_{21}R_{12}$, where P is a permutation matrix with the entry $P^{ij}_{kl} = \delta_{il}\delta_{jk}$. Then the braided-vectors algebra $V(R', R)$ defined by generators 1, $\{e^i \mid i = 1, \cdots, n\}$, and relations $e^i e^j = \sum_{a,b} R'^{\ ji}_{ab} e^a e^b$ forms a braided group with $\underline{\Delta}(e^i) = e^i \otimes 1 + 1 \otimes e^i$, $\underline{\epsilon}(e^i) = 0$, $\underline{S}(e^i) = -e^i$, $\Psi(e^i \otimes e^j) = \sum_{a,b} R^{ji}_{ab} e^a \otimes e^b$ in braided*

category ${}^{A(R)}\mathfrak{M}$. *Under the duality* $\langle f_j, e^i\rangle = \delta_{ij}$, *braided-covectors algebra* $V^\vee(R', R_{21}^{-1})$ *defined by* 1 *and* $\{f_j \mid j = 1, \cdots, n\}$, *and relations* $f_i f_j = \sum_{a,b} f_b f_a R'^{ab}_{ij}$ *forms another braided group with* $\underline{\Delta}(f_i) = f_i \otimes 1 + 1 \otimes f_i$, $\underline{\epsilon}(f_i) = 0$, $\underline{S}(f_i) = -f_i$, $\Psi(f_i \otimes f_j) = \sum_{a,b} f_b \otimes f_a R^{ab}_{ij}$ *in braided category* $\mathfrak{M}^{A(R)}$.

Remark 2.2. *There is a way to find suitable matrix* R' *for a given* R *in* [M4]. *We know that* PR *obeys some minimal polynomial* $\prod_i (PR - x_i) = 0$. *For each nonzero eigenvalue* x_i, *we can normalise* R *so that* $x_i = -1$. *then set* $R' = P + P\prod_{j\neq i}(PR - x_j)$, *which satisfies the conditions* (*i*)—(*iii*). *It gives us at least one braided (co)vector algebra for each nonzero eigenvalue of* PR.

With these, starting from some irreducible representation T_V of $U_q(\mathfrak{g})$ and its corresponding R-matrix R_{VV}. Then there exists a unique dual pairing $\langle\cdot,\cdot\rangle$ between $U_q^{\rm ext}(\mathfrak{g})$ and $H_{R_{VV}}$ in [HH2] such that

$$\langle (m^+)^i_j, t^k_l\rangle = R_{VV}{}^{ik}_{jl} = \lambda R^{ik}_{jl}, \quad \langle (m^-)^i_j, t^k_l\rangle = R^{-1\,ki}_{VV\,lj} = \lambda^{-1}(R^{-1})^{ki}_{lj},$$

$$\langle (m^+)^i_j, \tilde{t}^k_l\rangle = (R^{t_2}_{VV})^{-1\,ik}_{\ \ jl} = ((\lambda R)^{t_2})^{-1\,ik}_{\ \ jl}, \quad \langle (m^-)^i_j, \tilde{t}^k_l\rangle = [(R^{-1}_{VV})^{t_1}]^{-1\,lj}_{\ \ ki} = [(\lambda^{-1}R^{-1})^{t_1}]^{-1\,lj}_{\ \ ki}.$$

Such λ is called a *normalization constant of quantum groups*. The convolution-invertible algebra / anti-coalgebra maps $\mathscr{R}, \bar{\mathscr{R}}$ in $\hom(H_{\lambda R}, U_q^{\rm ext}(\mathfrak{g}))$ are

$$\mathscr{R}(t^i_j) = (m^+)^i_j, \quad \mathscr{R}(\tilde{t}^i_j) = (m^+)^{-1\,j}_{\ \ i}; \quad \bar{\mathscr{R}}(t^i_j) = (m^-)^i_j, \quad \bar{\mathscr{R}}(\tilde{t}^i_j) = (m^-)^{-1\,j}_{\ \ i},$$

where $U_q^{\rm ext}(\mathfrak{g})$ is the FRT-form of $U_q(\mathfrak{g})$ or extended quantized enveloping algebra adjoined by the elements $K_i^{\frac{1}{m}}$, $H_{\lambda R}$ is the Hopf algebra associated with $A(R_{VV})$, generated by $t^i_j, \tilde{t}^i_j$ and subject to certain relations. With the weakly quasitriangular dual pair of Hopf algebras in the above Theorem, we obtain that the braided (co)vector algebras $V(R', R) \in {}^{H_{\lambda R}}\mathfrak{M}$ and $V^\vee(R', R_{21}^{-1}) \in \mathfrak{M}^{H_{\lambda R}}$. However, in order to yield the required quantum group of higher rank 1, we need to centrally extend the pair $(U_q^{\rm ext}(\mathfrak{g}), H_{\lambda R})$ to the pair

$$\Big(\widetilde{U_q^{\rm ext}(\mathfrak{g})} = U_q^{\rm ext}(\mathfrak{g}) \otimes k[c, c^{-1}],\ \widetilde{H_{\lambda R}} = H_{\lambda R} \otimes k[g, g^{-1}]\Big)$$

with the action $e^i \triangleleft c = \lambda e^i$, $f_i \triangleleft c = \lambda f_i$ and the extended pair $\langle c, g\rangle = \lambda$. In this way, we easily see that the braided algebras $V(R', R) \in {}^{\widetilde{H_{\lambda R}}}\mathfrak{M}$ and $V^\vee(R', R_{21}^{-1}) \in \mathfrak{M}^{\widetilde{H_{\lambda R}}}$.

Having these, we have the following *generalized double-bosonization construction Theorem* in [HH2].

Theorem 2.2. *Let* R_{VV} *be the R-matrix corresponding to the irreducible representation* T_V *of* $U_q(\mathfrak{g})$. *There exists a normalization constant* λ *such that* $\lambda R = R_{VV}$. *Then the new*

quantum group $U = U(V^\vee(R', R_{21}^{-1}), \widetilde{U_q^{ext}(\mathfrak{g})}, V(R', R))$ *has the following the cross relations:*

$$cf_i = \lambda f_i c, \quad e^i c = \lambda c e^i, \quad [c, m^\pm] = 0, \quad [e^i, f_j] = \delta_{ij}\frac{(m^+)^i_j c^{-1} - c(m^-)^i_j}{q_* - q_*^{-1}};$$
$$e^i(m^+)^j_k = R_{VV}{}^{ji}_{ab}(m^+)^a_k e^b, \quad (m^-)^i_j e^k = R_{VV}{}^{ki}_{ab} e^a (m^-)^b_j,$$
$$(m^+)^i_j f_k = f_b (m^+)^i_a R_{VV}{}^{ab}_{jk}, \quad f_i (m^-)^j_k = (m^-)^j_b f_a R_{VV}{}^{ab}_{ik},$$

and the coproduct:

$$\Delta c = c \otimes c, \quad \Delta e^i = e^a \otimes (m^+)^i_a c^{-1} + 1 \otimes e^i, \quad \Delta f_i = f_i \otimes 1 + c(m^-)^a_i \otimes f_a,$$

and the counit $\epsilon e^i = \epsilon f_i = 0$. *We can normalize* e^i *such that the factor* $q_* - q_*^{-1}$ *meets the situation we need.*

Remark 2.3. *The R-matrix in* [M1] *is the* $P \circ \cdot \circ P$ *of the ordinary R-matrix, so the quantized enveloping algebras constructed by double-bosonization satisfy* $E_i K_j = q_i^{a_{ij}} K_j E_i$, $F_i K_j = q_i^{-a_{ij}} K_j F_i$, $[E_i, F_j] = \delta_{ij}\frac{K_i - K_i^{-1}}{q_i - q_i^{-1}}$, *where* $q_i = q^{d_i}$, *in addition to the same q-Serre relations. So it is necessary to notice that we use this definition of* $U_q(\mathfrak{g})$ *in our proof.*

Remark 2.4. *The R-matrix datum associated with a given module V comes from the* R_{VV}*-matrix, defined by* $R_{VV} = B_{VV} \circ (T_V \otimes T_V)(\mathfrak{R})$. *Here* $\mathfrak{R} = \sum_{r_1, \cdots, r_n = 0}^{\infty} \prod_{j=1}^{n} \frac{(1 - q_{\beta_j}^{-2})^{r_j}}{[r_j]_{q_{\beta_j}}} q_{\beta_j}^{\frac{r_j(r_j+1)}{2}} E_{\beta_j}^{r_j} \otimes F_{\beta_j}^{r_j}$ *is main part of the universal R-matrix of* $U_q(\mathfrak{g})$. B_{VV} *denotes the linear operator on* $V \otimes V$ *given by* $B_{VV}(v \otimes w) := q^{(\mu,\mu')} v \otimes w$ *for* $v \in V_\mu$, $w \in V_{\mu'}$. *We can endow every basis with weights indices such that the operators* $T_V(E_i)$ *and* $T_V(F_i)$ *are raising and lowering the indices of the basis, then the required* R_{VV}*-matrix is upper triangular (see* [HH2]*).*

Remark 2.5. *The matrix* $m^\pm$ *can be obtained by* $S(l_{ij}^\pm) = (m^\pm)^i_j$, *where* $l_{ij}^\pm$ *is the* $L^\pm$*-functionals of* $U_q(\mathfrak{g})$. *The* $L^\pm$*-functionals of* $U_q^{ext}(\mathfrak{g})$ *can obtained from* $U_h(\mathfrak{g})$*'s* $L^\pm$*-functionals in Chapter 8 in* [K], *defined by*

$$(id \otimes T_V)(\mathcal{R})(1 \otimes v_j) = \sum_i l_{ij}^+ \otimes v_i,$$
$$(T_V \otimes id)(\mathcal{R}^{-1})(v_j \otimes 1) = (T_V \otimes id)(S \otimes id)(\mathcal{R})(v_j \otimes 1) = \sum_i v_i \otimes l_{ij}^-,$$

where

$$\mathcal{R} = exp(h \sum_{i,j} B_{ij}(H_i \otimes H_j)) \prod_{\beta \in \Delta_+} exp_{q_\beta}((1 - q_\beta^{-2})(E_\beta \otimes F_\beta))$$

is the universal R-matrix of $U_h(\mathfrak{g})$. *Here matrix* (B_{ij}) *is the inverse of matrix* $(C_{ij}) = (d_j^{-1} a_{ij})$, *the q-exponential function* $exp_q x$ *is defined by* $exp_q x = \sum_{r=0}^{\infty} \frac{q^{\frac{r(r+1)}{2}}}{[r]_q!} x^r$.

3. Exceptional quantized enveloping algebras $U_q(\mathfrak{g})$ of type E series

From the type-crossing constructions of $U_q(G_2)$ and $U_q(F_4)$ in [HH2], we know that it is most crucial to find suitable representations, and get the R_{VV}-matrices we need, then to achieve the required data R, R' via determining the minimal polynomial of the PR_{VV}-matrix.

3.1. **Type-crossing construction of $U_q(E_6)$.** In order to construct $U_q(E_6)$, it is natural to start from $U_q(D_5)$ and a certain representation of it. We choose the 16-dimensional half-spin representation T_V of $U_q(D_5)$, which is given by the Figure 1, where $(s_1, s_2, s_3, s_4, s_5)$ denotes the basis of weight $\frac{1}{2}(s_1\varepsilon_1 + s_2\varepsilon_2 + s_3\varepsilon_3 + s_4\varepsilon_4 + s_5\varepsilon_5)$, $s_i = \pm$.



Figure 1. The 16-dimensional half-spin representation of $U_q(D_5)$

The actions of the E_i, F_i's can be read directly from the Figure 1, K_j's actions can be obtained by the defining relations of K_i and E_i, F_i in $U_q(D_5)$. Since $E_\beta(f_m) = f_n \Longleftrightarrow F_\beta(f_n) = f_m$ and E_i^2 is zero action, we can prove that the corresponding matrix PR_{VV} is symmetrizable, which is similar to situations in [HH2]. Moreover, for the corresponding R-matrix R_{VV}, we have

Lemma 3.1. (1) *The minimal polynomial of PR_{VV} is*

$$(PR_{VV} + q^{-\frac{3}{4}}I)(PR_{VV} - q^{\frac{5}{4}}I)(PR_{VV} - q^{-\frac{3}{4}}I) = 0.$$

(2) *Those entries we need in matrices $m^{\pm}$ are as follows:*

- *In m^+:*

$$(m^+)_2^1 = -(q-q^{-1})E_4K_1^{\frac{1}{2}}K_2K_3^{\frac{3}{2}}K_4^{\frac{1}{4}}K_5^{\frac{3}{4}}, \qquad (m^+)_2^2 = K_1^{\frac{1}{2}}K_2K_3^{\frac{3}{2}}K_4^{\frac{1}{4}}K_5^{\frac{3}{4}},$$
$$(m^+)_3^2 = -(q-q^{-1})E_3K_1^{\frac{1}{2}}K_2K_3^{\frac{1}{2}}K_4^{\frac{1}{4}}K_5^{\frac{3}{4}}, \qquad (m^+)_3^3 = K_1^{\frac{1}{2}}K_2K_3^{\frac{1}{2}}K_4^{\frac{1}{4}}K_5^{\frac{3}{4}},$$
$$(m^+)_4^3 = -(q-q^{-1})E_2K_1^{\frac{1}{2}}K_3^{\frac{1}{2}}K_4^{\frac{1}{4}}K_5^{\frac{3}{4}}, \qquad (m^+)_4^4 = K_1^{\frac{1}{2}}K_3^{\frac{1}{2}}K_4^{\frac{1}{4}}K_5^{\frac{3}{4}},$$
$$(m^+)_5^4 = -(q-q^{-1})E_1K_1^{-\frac{1}{2}}K_3^{\frac{1}{2}}K_4^{\frac{1}{4}}K_5^{\frac{3}{4}}, \qquad (m^+)_5^5 = K_1^{-\frac{1}{2}}K_3^{\frac{1}{2}}K_4^{\frac{1}{4}}K_5^{\frac{3}{4}},$$
$$(m^+)_{16}^{15} = -(q-q^{-1})E_5K_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{3}{4}}K_5^{-\frac{5}{4}}, \ (m^+)_{16}^{16} = K_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{3}{4}}K_5^{-\frac{5}{4}}.$$

- *In m^-:*

$$(m^-)_1^2 = q(q-q^{-1})K_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{1}{4}}K_5^{-\frac{3}{4}}F_4, \ (m^-)_2^2 = K_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{1}{4}}K_5^{-\frac{3}{4}},$$
$$(m^-)_{12}^{13} = q(q-q^{-1})K_1^{\frac{1}{2}}K_3^{\frac{1}{2}}K_4^{\frac{3}{4}}K_5^{\frac{1}{4}}F_1, \qquad (m^-)_{13}^{13} = K_1^{\frac{1}{2}}K_3^{\frac{1}{2}}K_4^{\frac{3}{4}}K_5^{\frac{1}{4}},$$
$$(m^-)_{13}^{14} = q(q-q^{-1})K_1^{\frac{1}{2}}K_2K_3^{\frac{1}{2}}K_4^{\frac{3}{4}}K_5^{\frac{1}{4}}F_2, \qquad (m^-)_{14}^{14} = K_1^{\frac{1}{2}}K_2K_3^{\frac{1}{2}}K_4^{\frac{3}{4}}K_5^{\frac{1}{4}},$$
$$(m^-)_{14}^{15} = q(q-q^{-1})K_1^{\frac{1}{2}}K_2K_3^{\frac{3}{2}}K_4^{\frac{3}{4}}K_5^{\frac{1}{4}}F_3, \qquad (m^-)_{15}^{15} = K_1^{\frac{1}{2}}K_2K_3^{\frac{3}{2}}K_4^{\frac{3}{4}}K_5^{\frac{1}{4}},$$
$$(m^-)_{15}^{16} = q(q-q^{-1})K_1^{\frac{1}{2}}K_2K_3^{\frac{3}{2}}K_4^{\frac{3}{4}}K_5^{\frac{5}{4}}F_5, \qquad (m^-)_{16}^{16} = K_1^{\frac{1}{2}}K_2K_3^{\frac{3}{2}}K_4^{\frac{3}{4}}K_5^{\frac{5}{4}}.$$

Proof. Those entries in (2) of Lemma 3.1 can be obtained by Remark 2.5. Then we will focus on the minimal polynomial in (1). The decomposition of tensor product for this spin representation is $T_V \otimes T_V = V_1 \oplus V_2 \oplus V_3$, where $V_i, i = 1, 2, 3$ is the irreducible representation with highest weight $\varepsilon_1 + \varepsilon_2 + \varepsilon_3 + \varepsilon_4 + \varepsilon_5, \varepsilon_1 + \varepsilon_2 + \varepsilon_3, \varepsilon_1$, respectively. This means that matrix PR_{VV} has only 3 distinct eigenvalues, denoted by x_1, x_2, x_3. Then PR_{VV} obeys the minimal polynomial $(PR_{VV} - x_1I)(PR_{VV} - x_2I)(PR_{VV} - x_3I) = 0$. We use an ingenious method to get these eigenvalues x_i.

According to Remark 2.4, we obtain the nonzero entries of $PR_{VV} - x_iI$ in rows of (12), (21) are

$$(PR_{VV} - x_iI)_{12}^{12} = -x_i,$$
$$(PR_{VV} - x_iI)_{21}^{12} = q^{\frac{1}{4}},$$
$$(PR_{VV} - x_iI)_{12}^{21} = q^{\frac{1}{4}},$$
$$(PR_{VV} - x_iI)_{21}^{21} = q^{\frac{1}{4}}(q - q^{-1}) - x_i.$$

Then we obtain that the nonzero entries in row (12) of $(PR_{VV} - x_1I)(PR_{VV} - x_2I)$ still lie in columns (12) and (21), given by

$$[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]_{12}^{12} = (PR_{VV}-x_1I)_{12}^{12}(PR_{VV}-x_2I)_{12}^{12}+(PR_{VV}-x_1I)_{21}^{12}(PR_{VV}-x_2I)_{12}^{21},$$
$$[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]_{21}^{12} = (PR_{VV}-x_1I)_{12}^{12}(PR_{VV}-x_2I)_{21}^{12}+(PR_{VV}-x_1I)_{21}^{12}(PR_{VV}-x_2I)_{21}^{21}.$$

With these, some entries in row (12) of $(PR_{VV}-x_1I)(PR_{VV}-x_2I)(PR_{VV}-x_3I)$ can be obtained by

$$\begin{aligned}
&[(PR_{VV}-x_1I)(PR_{VV}-x_2I)(PR_{VV}-x_3I)]^{12}_{12}\\
&\quad=[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]^{12}_{12}(PR_{VV}-x_3I)^{12}_{12}\\
&\qquad+[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]^{12}_{21}(PR_{VV}-x_3I)^{21}_{12}\\
&\quad=-x_1x_2x_3-q^{\frac{1}{2}}(x_1+x_2+x_3)+q^{\frac{3}{4}}(q-q^{-1}),\\
&[(PR_{VV}-x_1I)(PR_{VV}-x_2I)(PR_{VV}-x_3I)]^{12}_{21}\\
&\quad=[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]^{12}_{12}(PR_{VV}-x_3I)^{12}_{21}\\
&\qquad+[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]^{12}_{21}(PR_{VV}-x_3I)^{21}_{21}\\
&\quad=q^{\frac{1}{4}}(x_1x_2+x_2x_3+x_1x_3)-q^{\frac{1}{2}}(q-q^{-1})(x_1+x_2+x_3)+q^{\frac{3}{4}}(q-q^{-1})^2+q^{\frac{3}{4}}.
\end{aligned}$$

We consider the nonzero entries in rows $(1,16),(16,1)$ as follows

$$\begin{aligned}
(PR_{VV}-x_iI)^{1,16}_{1,16}&=-x_i,\\
(PR_{VV}-x_iI)^{1,\ 16}_{16,1}&=q^{-\frac{3}{4}},\\
(PR_{VV}-x_iI)^{16,1}_{16,1}&=-x_i,\\
(PR_{VV}-x_iI)^{16,1}_{1,\ 16}&=q^{-\frac{3}{4}},\\
(PR_{VV}-x_iI)^{16,1}_{a_0,b_0}&=Y_{a_0}
\end{aligned}$$

for some columns (a_0,b_0). If there exists $Y_{a_0}\neq 0$, then $1<a_0,\ b_0<16$, so

$$(PR_{VV}-x_iI)^{a_0,b_0}_{1,\ 16}=0.$$

Nonzero entries in row $(1,16)$ of $(PR_{VV}-x_1I)(PR_{VV}-x_2I)$ are given by

$$\begin{aligned}
&[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]^{1,16}_{1,16}\\
&\quad=(PR_{VV}-x_1I)^{1,16}_{1,16}(PR_{VV}-x_2I)^{1,16}_{1,16}+(PR_{VV}-x_1I)^{1,\ 16}_{16,1}(PR_{VV}-x_2I)^{16,1}_{1,\ 16},\\
&[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]^{1,\ 16}_{16,1}\\
&\quad=(PR_{VV}-x_1I)^{1,16}_{1,16}(PR_{VV}-x_2I)^{1,\ 16}_{16,1}+(PR_{VV}-x_1I)^{1,\ 16}_{16,1}(PR_{VV}-x_2I)^{16,1}_{16,1},\\
&[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]^{1,\ 16}_{a_0,b_0}\\
&\quad=(PR_{VV}-x_1I)^{1,16}_{1,16}(PR_{VV}-x_2I)^{1,\ 16}_{a_0,b_0}+(PR_{VV}-x_1I)^{1,\ 16}_{16,1}(PR_{VV}-x_2I)^{16,1}_{a_0,b_0}\\
&\quad=(PR_{VV}-x_1I)^{1,\ 16}_{16,1}(PR_{VV}-x_2I)^{16,1}_{a_0,b_0}.
\end{aligned}$$

With these, the entry $[(PR_{VV}-x_1I)(PR_{VV}-x_2I)(PR_{VV}-x_3I)]^{1,16}_{1,16}$ is

$$\begin{aligned}
&[(PR_{VV}-x_1I)(PR_{VV}-x_2I)(PR_{VV}-x_3I)]^{1,16}_{1,16}\\
&\quad=[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]^{1,16}_{1,16}(PR_{VV}-x_3I)^{1,16}_{1,16}\\
&\qquad+[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]^{1,\ 16}_{16,1}(PR_{VV}-x_3I)^{16,1}_{1,\ 16}\\
&\qquad+[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]^{1,\ 16}_{a_0,b_0}(PR_{VV}-x_3I)^{a_0,b_0}_{1,\ 16}\\
&\quad=[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]^{1,16}_{1,16}(PR_{VV}-x_3I)^{1,16}_{1,16}\\
&\qquad+[(PR_{VV}-x_1I)(PR_{VV}-x_2I)]^{1,\ 16}_{16,1}(PR_{VV}-x_3I)^{16,1}_{1,\ 16}\\
&\quad=-x_1x_2x_3-q^{-\frac{3}{2}}(x_1+x_2+x_3).
\end{aligned}$$

Then we have the following equations

$$\begin{cases}
-x_1x_2x_3-q^{\frac{1}{2}}(x_1+x_2+x_3)+q^{\frac{3}{4}}(q-q^{-1})=0,\\
q^{\frac{1}{4}}(x_1x_2+x_2x_3+x_1x_3)-q^{\frac{1}{2}}(q-q^{-1})(x_1+x_2+x_3)+q^{\frac{3}{4}}(q-q^{-1})^2+q^{\frac{3}{4}}=0,\\
-x_1x_2x_3-q^{-\frac{3}{2}}(x_1+x_2+x_3)=0.
\end{cases}$$

Solving these equations, we obtain that these 3 eigenvalues are $q^{\frac{5}{4}}, q^{-\frac{3}{4}}, -q^{-\frac{3}{4}}$. So the minimal polynomial of PR_{VV} is $(PR_{VV}+q^{-\frac{3}{4}}I)(PR_{VV}-q^{\frac{5}{4}}I)(PR_{VV}-q^{-\frac{3}{4}}I)=0$. □

From the above Lemma 3.1, setting

$$R=q^{\frac{3}{4}}R_{VV},\quad R'=RPR-(q^2+1)R+(q^2+1)P,$$

then we obtain $(PR+I)(PR'-I)=0$. With these, we have the following

Theorem 3.1. *With the quantum normalization constant $\lambda=q^{-\frac{3}{4}}$, identify e^{16}, f_{16}, and $(m^+)^{16}_{16}c^{-1}$ with the additional simple root vectors E_6, F_6, and the group-like element K_6, respectively, then the new quantum group $U(V^\vee(R',R_{21}^{-1}), U_q^{\widetilde{ext}}(D_5), V(R',R))$ is the quantized enveloping algebra $U_q(E_6)$ with $K_i^{\frac{1}{4}}$ adjoined.*

Proof. Under the above identification, $[E_6,F_6]=\frac{K_6-K_6^{-1}}{q-q_{-1}}$, $\Delta(E_6)=E_6\otimes K_6+1\otimes E_6$, and $\Delta(F_6)=F_6\otimes 1+K_6^{-1}\otimes F_6$ can be obtained directly by Theorem 2.2. We find that the minor diagonal entries $(m^-)^{i+1}_i$ has the form $(m^-)^{i+1}_i=q(q-q^{-1})(m^-)^{i+1}_{i+1}F$. Combing with $(m^-)^{i+1}_ie^{16}=\lambda R^{16i+1}_{16i+1}e^{16}(m^-)^{i+1}_i$ and $(m^-)^{i+1}_{i+1}e^{16}=\lambda R^{16i+1}_{16i+1}e^{16}(m^-)^{i+1}_{i+1}$ given by the equalities in Theorem 2.2, we obtain $[E_6,F_j]=0, 1\le j\le 5$, namely, $[E_6,F_i]=\delta_{ij}\frac{K_6-K_6^{-1}}{q-q_{-1}}$. According to the cross relations in Theorem 2.2 again, we have

$$\left.\begin{array}{l}
e^{16}(m^+)^j_j=\lambda R^{j\ j}_{1616}(m^+)^j_je^{16},\\
(m^+)^{i+1}_{i+1}K_{5-i}=(m^+)^i_i,\quad 1\le i\le 4,\\
R^{i\ \ i}_{1616}=1,\quad 1\le i\le 5,\quad R^{1616}_{1616}=q^2
\end{array}\right\}\Longrightarrow\begin{cases}
E_6K_i=K_iE_6,\quad 1\le i\le 4,\\
E_6K_5=q^{-1}K_5E_6,\\
E_6K_6=q^2K_6E_6.
\end{cases}$$

The relations between K_6 and others E_i's are given by

$$E_1K_6 = E_1K_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{3}{4}}K_5^{-\frac{5}{4}}c^{-1} = q^{-1}qK_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{3}{4}}K_5^{-\frac{5}{4}}c^{-1}E_1 = K_6E_1.$$
$$E_2K_6 = E_2K_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{3}{4}}K_5^{-\frac{5}{4}}c^{-1} = q^{\frac{1}{2}}q^{-2}q^{\frac{3}{2}}K_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{3}{4}}K_5^{-\frac{5}{4}}c^{-1}E_2 = K_6E_2,$$
$$E_3K_6 = E_3K_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{3}{4}}K_5^{-\frac{5}{4}}c^{-1} = qq^{-3}q^{\frac{3}{4}}q^{\frac{5}{4}}K_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{3}{4}}K_5^{-\frac{5}{4}}c^{-1}E_3 = K_6E_3,$$
$$E_4K_6 = E_4K_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{3}{4}}K_5^{-\frac{5}{4}}c^{-1} = q^{\frac{1}{2}}q^{-2}q^{\frac{3}{2}}K_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{3}{4}}K_5^{-\frac{5}{4}}c^{-1}E_4 = K_6E_4,$$
$$E_5K_6 = E_5K_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{3}{4}}K_5^{-\frac{5}{4}}c^{-1} = q^{\frac{3}{2}}q^{-\frac{5}{2}}K_1^{-\frac{1}{2}}K_2^{-1}K_3^{-\frac{3}{2}}K_4^{-\frac{3}{4}}K_5^{-\frac{5}{4}}c^{-1}E_5 = q^{-1}K_6E_5.$$

We will explore the q-Serre relations between E_6 and others E_i's. E_i's appear in the minor entries of m^+, so we consider the cross relations related to $(m^+)^i_{i+1}$ in Theorem 2.2.

$$\left.\begin{array}{l}(m^+)^i_{i+1} = -(q-q^{-1})E_{5-i}(m^+)^{i+1}_{i+1},\\ e^{16}(m^+)^i_{i+1} = \lambda(m^+)^i_{i+1}e^{16},\\ e^{16}(m^+)^{i+1}_{i+1} = \lambda(m^+)^{i+1}_{i+1}e^{16}, 1\le i\le 4,\\ e^{16}(m^+)^{15}_{16} = \lambda(q(m^+)^{15}_{16}e^{16} + q(q-q^{-1})(m^+)^{16}_{16}e^{15},\end{array}\right\} \Longrightarrow \left\{\begin{array}{l}E_6E_i - E_iE_6 = 0,\\ e^{15} = qe^{16}E_5 - E_5e^{16}.\end{array}\right.$$

So we need to know the relations between e^{15} and e^{16}, E_5. Indeed, $e^{15}e^{16} = R'^{16,15}_{a,\ b}e^ae^b = (2q^2+1)e^{15}e^{16} - 2qe^{16}e^{15} \Longrightarrow e^{16}e^{15} = qe^{15}e^{16}$, then combing with the above equality $e^{15} = qe^{16}E_5 - E_5e^{16}$, we obtain

$$(E_6)^2E_5 - [2]_qE_6E_5E_6 + E_5(E_6)^2 = 0.$$

On the other hand, the relation of e^{15} and E_5 is given by $e^{15}(m^+)^{15}_{16} = \lambda R^{1515}_{a\ b}(m^+)^a_{16}e^b = q^{\frac{5}{4}}(m^+)^{15}_{16}e^{15}$. Putting the expression of $(m^+)^{15}_{16}$ into the equality, we get $e^{15}E_5 = qE_5e^{15}$. Combing with the above equality $e^{15} = qe^{16}E_5 - E_5e^{16}$ again, we obtain

$$(E_5)^2E_6 - [2]_qE_5E_6E_5 + E_6(E_5)^2 = 0.$$

The relations of f_{16} and F_i, K_i can be obtained in a similar way. □

3.2. **Inductive construction of $U_q(E_7)$.** For the construction of $U_q(E_7)$, we choose the 27-dimensional minimal fundamental representation of $U_q(E_6)$ given by the following Figure 2, which also appeared in [KM, KKM]. $(s_1, \cdots s_i)$ denotes the basis of weight $s_1\lambda_1 + \cdots + s_i\lambda_i$, $s_j = 0, 1, \overline{1}$, where $\overline{1} = -1$. Corresponding to this representation, we have the following Lemma 3.2 by the similar analysis.

Lemma 3.2. (1) *PR_{VV} obeys the minimal polynomial*

$$(PR_{VV} - q^{\frac{4}{3}}I)(PR_{VV} - q^{-\frac{2}{3}}I)(PR_{VV} + q^{-\frac{2}{3}}I) = 0.$$

(2) *The entries we need in $m^{\pm}$ are obtained by Remark 2.5,*

- *In* m^+ :

$$(m^+)_2^1 = (q^{-1}-q)E_1K_1^{\frac{1}{3}}K_2K_3^{\frac{5}{3}}K_4^2K_5^{\frac{4}{3}}K_6^{\frac{2}{3}}, \quad (m^+)_2^2 = K_1^{\frac{1}{3}}K_2K_3^{\frac{5}{3}}K_4^2K_5^{\frac{4}{3}}K_6^{\frac{2}{3}},$$
$$(m^+)_3^2 = (q^{-1}-q)E_3K_1^{\frac{1}{3}}K_2K_3^{\frac{2}{3}}K_4^2K_5^{\frac{4}{3}}K_6^{\frac{2}{3}}, \quad (m^+)_3^3 = K_1^{\frac{1}{3}}K_2K_3^{\frac{2}{3}}K_4^2K_5^{\frac{4}{3}}K_6^{\frac{2}{3}},$$
$$(m^+)_4^3 = (q^{-1}-q)E_4K_1^{\frac{1}{3}}K_2K_3^{\frac{2}{3}}K_4K_5^{\frac{4}{3}}K_6^{\frac{2}{3}}, \quad (m^+)_4^4 = K_1^{\frac{1}{3}}K_2K_3^{\frac{2}{3}}K_4K_5^{\frac{4}{3}}K_6^{\frac{2}{3}},$$
$$(m^+)_5^4 = (q^{-1}-q)E_2K_1^{\frac{1}{3}}K_3^{\frac{2}{3}}K_4K_5^{\frac{4}{3}}K_6^{\frac{2}{3}}, \quad (m^+)_5^5 = K_1^{\frac{1}{3}}K_3^{\frac{2}{3}}K_4K_5^{\frac{4}{3}}K_6^{\frac{2}{3}}, \quad (m^+)_6^5 = 0,$$
$$(m^+)_6^4 = (q^{-1}-q)E_5K_1^{\frac{1}{3}}K_2K_3^{\frac{2}{3}}K_4K_5^{\frac{1}{3}}K_6^{\frac{2}{3}}, \quad (m^+)_6^6 = K_1^{\frac{1}{3}}K_2K_3^{\frac{2}{3}}K_4K_5^{\frac{1}{3}}K_6^{\frac{2}{3}},$$
$$(m^+)_7^6 = (q^{-1}-q)E_6K_1^{\frac{1}{3}}K_2K_3^{\frac{2}{3}}K_4K_5^{\frac{1}{3}}K_6^{-\frac{1}{3}}, \quad (m^+)_7^7 = K_1^{\frac{1}{3}}K_2K_3^{\frac{2}{3}}K_4K_5^{\frac{1}{3}}K_6^{-\frac{1}{3}},$$
$$(m^+)_1^1 = K_1^{\frac{4}{3}}K_2K_3^{\frac{5}{3}}K_4^2K_5^{\frac{4}{3}}K_6^{\frac{2}{3}}, \quad (m^+)_{27}^{27} = K_1^{-\frac{2}{3}}K_2^{-1}K_3^{-\frac{4}{3}}K_4^{-2}K_5^{-\frac{5}{3}}K_6^{-\frac{4}{3}}.$$

- *In* m^- :

$$(m^-)_1^2 = \frac{K_1^{-\frac{1}{3}}K_2^{-1}K_3^{-\frac{5}{3}}K_4^{-2}K_5^{-\frac{4}{3}}K_6^{-\frac{2}{3}}F_1}{(q^2-1)}, \quad (m^-)_2^2 = K_1^{-\frac{1}{3}}K_2^{-1}K_3^{-\frac{5}{3}}K_4^{-2}K_5^{-\frac{4}{3}}K_6^{-\frac{2}{3}},$$
$$(m^-)_2^3 = \frac{K_1^{-\frac{1}{3}}K_2^{-1}K_3^{-\frac{2}{3}}K_4^{-2}K_5^{-\frac{4}{3}}K_6^{-\frac{2}{3}}F_3}{(q^2-1)}, \quad (m^-)_3^3 = K_1^{-\frac{1}{3}}K_2^{-1}K_3^{-\frac{2}{3}}K_4^{-2}K_5^{-\frac{4}{3}}K_6^{-\frac{2}{3}},$$
$$(m^-)_3^4 = \frac{K_1^{-\frac{1}{3}}K_2^{-1}K_3^{-\frac{2}{3}}K_4^{-1}K_5^{-\frac{4}{3}}K_6^{-\frac{2}{3}}F_4}{(q^2-1)}, \quad (m^-)_4^4 = K_1^{-\frac{1}{3}}K_2^{-1}K_3^{-\frac{2}{3}}K_4^{-1}K_5^{-\frac{4}{3}}K_6^{-\frac{2}{3}},$$
$$(m^-)_4^5 = \frac{K_1^{-\frac{1}{3}}K_3^{-\frac{2}{3}}K_4^{-1}K_5^{-\frac{4}{3}}K_6^{-\frac{2}{3}}F_2}{(q^2-1)}, \quad (m^-)_5^5 = K_1^{-\frac{1}{3}}K_3^{-\frac{2}{3}}K_4^{-1}K_5^{-\frac{4}{3}}K_6^{-\frac{2}{3}},$$
$$(m^-)_5^8 = \frac{K_1^{-\frac{1}{3}}K_3^{-\frac{2}{3}}K_4^{-1}K_5^{-\frac{1}{3}}K_6^{-\frac{2}{3}}F_5}{(q^2-1)}, \quad (m^-)_8^8 = K_1^{-\frac{1}{3}}K_3^{-\frac{2}{3}}K_4^{-1}K_5^{-\frac{1}{3}}K_6^{-\frac{2}{3}},$$
$$(m^-)_6^7 = \frac{K_1^{-\frac{1}{3}}K_2^{-1}K_3^{-\frac{2}{3}}K_4^{-1}K_5^{-\frac{1}{3}}K_6^{\frac{1}{3}}F_6}{(q^2-1)}, \quad (m^-)_7^7 = K_1^{-\frac{1}{3}}K_2^{-1}K_3^{-\frac{2}{3}}K_4^{-1}K_5^{-\frac{1}{3}}K_6^{\frac{1}{3}}.$$

Proof. The method of proof is similar to the above Lemma. For the proof of (1), the decomposition of tensor product for this representation is

$$(000001) \otimes (000001) = (000002) \oplus (000010) \oplus (100000).$$

Namely, $V \otimes V = V_1 \oplus V_2 \oplus V_3$, where $V_i, i = 1, 2, 3$ denotes the irreducible representation with highest weight $2\lambda_6, \lambda_5, \lambda_1$, respectively. This means that matrix PR_{VV} has 3 distinct eigenvalues, denoted by x_1, x_2, x_3, and set $\mathcal{N} = (PR_{VV} - x_1I)(PR_{VV} - x_2I)(PR_{VV} - x_3I)$.

We also consider some entries at some special rows in the matrix $\mathcal{N}$. For example, the nonzero entries at rows (12) and (21) are $(PR_{VV} - x_iI)_{12}^{12} = -x_i, (PR_{VV} - x_iI)_{21}^{12} = q^{\frac{1}{3}}, (PR_{VV} - x_iI)_{12}^{21} = q^{\frac{1}{3}}, (PR_{VV} - x_iI)_{21}^{21} = q^{\frac{1}{3}}(q - q^{-1}) - x_i$. Then we obtain

$$\mathcal{N}_{12}^{12} = -x_1x_2x_3 - q^{\frac{2}{3}}(x_1 + x_2 + x_3) + q^2 - 1,$$
$$\mathcal{N}_{21}^{12} = q^{\frac{1}{3}}(x_1x_2 + x_1x_3 + x_2x_3) - q^{\frac{2}{3}}(q - q^{-1})(x_1 + x_2 + x_3) + q^3 + q^{-1} - q.$$

Other special rows we consider are $(1, 27)$ and $(27, 1)$, nonzero entries at these rows are $(PR_{VV} - x_iI)_{1,27}^{1,27} = -x_i, (PR_{VV} - x_iI)_{27,1}^{1,\ 27} = q^{-\frac{3}{4}}, (PR_{VV} - x_iI)_{27,1}^{27,1} = -x_i, (PR_{VV} - x_iI)_{1,\ 27}^{27,1} = q^{-\frac{3}{4}}, (PR_{VV} - x_iI)_{c_0,\,d_0}^{27,1} = Y_{c_0}$ for some columns (a_0, b_0), where $1 < a_0, b_0 < 27$,

and $(PR_{VV} - x_i I)^{a_0,b_0}_{1,\ 27} = 0$. With these entries, we obtain

$$\mathcal{N}^{1,27}_{1,27} = -x_1x_2x_3 - q^{-\frac{4}{3}}(x_1 + x_2 + x_3).$$

In view of the value of $\mathcal{N}^{12}_{12}, \mathcal{N}^{12}_{21}, \mathcal{N}^{1,27}_{1,27}$, we obtain the following equations

$$\begin{cases} -x_1x_2x_3 - q^{\frac{2}{3}}(x_1 + x_2 + x_3) + q^2 - 1 = 0, \\ q^{\frac{1}{3}}(x_1x_2 + x_1x_3 + x_2x_3) - q^{\frac{2}{3}}(q - q^{-1})(x_1 + x_2 + x_3) + q^3 + q^{-1} - q = 0, \\ -x_1x_2x_3 - q^{-\frac{4}{3}}(x_1 + x_2 + x_3) = 0. \end{cases}$$

Solving it, we get that these eigenvalues are $q^{\frac{4}{3}}, \pm q^{-\frac{2}{3}}$. Then we obtain the minimal polynomial of PR_{VV} is $(PR_{VV} - q^{\frac{4}{3}}I)(PR_{VV} - q^{-\frac{2}{3}}I)(PR_{VV} + q^{-\frac{2}{3}}I) = 0$. □

Setting $R = q^{\frac{2}{3}}R_{VV}, R' = RPR - (q^2 + 1)R + (q^2 + 1)P$, we have $(PR + I)(PR' - I) = 0$. In view of these results, we obtain the following

Theorem 3.2. *With the quantum normalization constant $\lambda = q^{-\frac{2}{3}}$, identify e^{27}, f_{27} and $(m^+)^{27}_{27}c^{-1}$ with the additional simple root vectors E_7, F_7 and the group-element K_7, respectively, then the new quantum group $U(V^\vee(R', R^{-1}_{21}), \widetilde{U^{ext}_q(E_6)}, V(R', R))$ is the quantized enveloping algebra $U_q(E_7)$ with $K_i^{\frac{1}{3}}$ adjoined.*

Proof. $[E_7, F_7] = \frac{K_7 - K_7^{-1}}{q - q_{-1}}$, $\Delta(E_7) = E_7 \otimes K_7 + 1 \otimes E_7$, and $\Delta(F_7) = F_7 \otimes 1 + K_7^{-1} \otimes F_7$ can by obtained directly by Theorem 2.2. We will explore the relations between E_7 and F_i. F_i's appear in the minor entries $(m^-)^{i+1}_i$ of matrix m^-, then according to the cross relations in Theorem 2.2, we obtain

$$\left.\begin{array}{l} (m^-)^{i+1}_i = (q^2 - 1)(m^-)^{i+1}_{i+1}F_j, \\ (m^-)^8_5 = (q^2 - 1)(m^-)^8_8F_5, \\ (m^-)^{i+1}_i e^{27} = \lambda R^{27\,i+1}_{27\,i+1} e^{27}(m^-)^{i+1}_i, \\ (m^-)^8_5 e^{27} = \lambda R^{278}_{278} e^{27}(m^-)^8_5, \\ (m^-)^{i+1}_{i+1} e^{27} = \lambda R^{27\,i+1}_{27\,i+1} e^{27}(m^-)^{i+1}_{i+1}, \end{array}\right\} \Longrightarrow E_7F_j = F_jE_7 \Longrightarrow [E_7, F_j] = 0, \quad 1 \le j \le 6.$$

Namely, $[E_7, F_i] = \delta_{7j}\frac{K_7 - K_7^{-1}}{q - q_{-1}}$. Under the above identification, $E_7K_7 = e^{27}(m^+)^{27}_{27}c^{-1} = \lambda R^{2727}_{2727}(m^+)^{27}_{27}e^{27}c^{-1} = R^{2727}_{2727}(m^+)^{27}_{27}c^{-1}e^{27} = q^{\frac{2}{3}}R_{VV}{}^{2727}_{2727}(m^+)^{27}_{27}c^{-1}e^{27} = q^2(m^+)^{27}_{27}c^{-1}e^{27} = q^2K_7E_7$. According to the cross relation $e^{27}(m^+)^i_i = \lambda R^{i27}_{i27}(m^+)^i_i e^{27}$ in Theorem 2.2, combining with $(m^+)^2_2K_1 = (m^+)^1_1$, $(m^+)^3_3K_3 = (m^+)^2_2$, $(m^+)^4_4K_4 = (m^+)^3_3$, $(m^+)^5_5K_2 = (m^+)^4_4$, $(m^+)^7_7K_6 = (m^+)^5_5$, and $R^{i27}_{i27} = 1, 1 \le i \le 6$, $R^{727}_{727} = q$, we obtain the relations between E_7 and K_i: $\begin{cases} E_7K_i = K_iE_7, \ 1 \le i \le 5, \\ E_7K_6 = q^{-1}K_6E_7. \end{cases}$

The relations between K_7 and others E_i's can be obtained by the following equalities.

$$E_1K_7 = E_1K_1^{-\frac{2}{3}}K_2^{-1}K_3^{-\frac{4}{3}}K_4^{-2}K_5^{-\frac{5}{3}}K_6^{-\frac{4}{3}}c^{-1} = q^{-\frac{4}{3}}q^{\frac{4}{3}}K_1^{-\frac{2}{3}}K_2^{-1}K_3^{-\frac{4}{3}}K_4^{-2}K_5^{-\frac{5}{3}}K_6^{-\frac{4}{3}}c^{-1}E_1 = K_7E_1,$$

$$E_2K_7 = E_2K_1^{-\frac{2}{3}}K_2^{-1}K_3^{-\frac{4}{3}}K_4^{-2}K_5^{-\frac{5}{3}}K_6^{-\frac{4}{3}}c^{-1} = q^{-2}q^2K_1^{-\frac{2}{3}}K_2^{-1}K_3^{-\frac{4}{3}}K_4^{-2}K_5^{-\frac{5}{3}}K_6^{-\frac{4}{3}}c^{-1}E_2 = K_7E_2,$$
$$E_3K_7 = E_3K_1^{-\frac{2}{3}}K_2^{-1}K_3^{-\frac{4}{3}}K_4^{-2}K_5^{-\frac{5}{3}}K_6^{-\frac{4}{3}}c^{-1} = q^{\frac{2}{3}}q^{-\frac{8}{3}}q^2K_1^{-\frac{2}{3}}K_2^{-1}K_3^{-\frac{4}{3}}K_4^{-2}K_5^{-\frac{5}{3}}K_6^{-\frac{4}{3}}c^{-1}E_3 = K_7E_3,$$
$$E_4K_7 = E_4K_1^{-\frac{2}{3}}K_2^{-1}K_3^{-\frac{4}{3}}K_4^{-2}K_5^{-\frac{5}{3}}K_6^{-\frac{4}{3}}c^{-1} = qq^{\frac{4}{3}}q^{-4}q^{\frac{5}{3}}K_1^{-\frac{2}{3}}K_2^{-1}K_3^{-\frac{4}{3}}K_4^{-2}K_5^{-\frac{5}{3}}K_6^{-\frac{4}{3}}c^{-1}E_4 = K_7E_4,$$
$$E_5K_7 = E_5K_1^{-\frac{2}{3}}K_2^{-1}K_3^{-\frac{4}{3}}K_4^{-2}K_5^{-\frac{5}{3}}K_6^{-\frac{4}{3}}c^{-1} = q^2q^{-\frac{10}{3}}q^{\frac{4}{3}}K_1^{-\frac{2}{3}}K_2^{-1}K_3^{-\frac{4}{3}}K_4^{-2}K_5^{-\frac{5}{3}}K_6^{-\frac{4}{3}}c^{-1}E_5 = K_7E_5,$$
$$E_6K_7 = E_6K_1^{-\frac{2}{3}}K_2^{-1}K_3^{-\frac{4}{3}}K_4^{-2}K_5^{-\frac{5}{3}}K_6^{-\frac{4}{3}}c^{-1} = q^{\frac{5}{3}}q^{-\frac{8}{3}}K_1^{-\frac{2}{3}}K_2^{-1}K_3^{-\frac{4}{3}}K_4^{-2}K_5^{-\frac{5}{3}}K_6^{-\frac{4}{3}}c^{-1}E_5 = q^{-1}K_7E_6.$$

We will describe the q-Serre relations between e^{27} and E_i. Since E_i's appear in the minor entries $(m^+)^i_{i+1}$ of matrix m^+, we have

$$\left.\begin{array}{l}(m^+)^i_{i+1} = -(q-q^{-1})E_j(m^+)^{i+1}_{i+1},\\ e^{27}(m^+)^i_{i+1} = \lambda R^{i27}_{i27}(m^+)^i_{i+1}e^{27}, i = 1,\cdots,4,\\ e^{27}(m^+)^{i+1}_{i+1} = \lambda R^{i+127}_{i+127}(m^+)^{i+1}_{i+1}e^{27},\end{array}\right\} \Longrightarrow\ E_7E_j = E_jE_7,\ j = 1,2,3,4.$$

$$\left.\begin{array}{l}(m^+)^4_6 = -(q-q^{-1})E_5(m^+)^6_6,\\ e^{27}(m^+)^4_6 = \lambda R^{427}_{ab}(m^+)^a_6e^b,\\ e^{27}(m^+)^6_6 = \lambda R^{627}_{627}(m^+)^6_6e^{27},\\ (m^+)^5_6 = 0, R^{427}_{625} = 0, R^{427}_{427} = R^{627}_{627} = 1,\end{array}\right\} \Longrightarrow\ E_7E_5 = E_5E_7.$$

$$\left.\begin{array}{l}(m^+)^6_7 = -(q-q^{-1})E_6(m^+)^7_7,\\ e^{27}(m^+)^6_7 = \lambda(m^+)^6_7e^{27} + \lambda(q-q^{-1})(m^+)^7_7e^{26},\\ e^{27}(m^+)^7_7 = \lambda R^{727}_{727}(m^+)^7_7e^{27} = \lambda q(m^+)^7_7e^{27},\\ e^{26}(m^+)^7_7 = \lambda R^{726}_{726}(m^+)^7_7e^{27} = \lambda(m^+)^7_7e^{26},\end{array}\right\} \Longrightarrow e^{26} = q^{-1}E_6e^{27} - e^{27}E_6.$$

According to $R'^{27,26}_{27,26} = -2q$, and $R'^{27,26}_{26,27} = 2q^2+1$, then $e^{26}e^{27} = R'^{27,\ 26}_{a,\ b}e^ae^b = -2qe^{27}e^{26} + (2q^2+1)e^{26}e^{27} \Longrightarrow e^{27}e^{26} = qe^{26}e^{27}$. Combing with the above relation $e^{26} = q^{-1}E_6e^{27} - e^{27}E_6$, we obtain

$$(E_7)^2E_6 - (q+q^{-1})E_7E_6E_7 + E_6(E_7)^2 = 0.$$

On the other hand, we need to know the relation between e^{26} and E_6, which can be explored by the following relations

$$\left\{\begin{array}{l}(m^+)^6_7 = (q-q^{-1})E_6(m^+)^7_7,\\ e^{26}(m^+)^6_7 = \lambda R^{626}_{626}(m^+)^6_7e^{26} = \lambda q(m^+)^6_7e^{26},\\ e^{26}(m^+)^7_7 = \lambda R^{726}_{726}(m^+)^7_7e^{26} = \lambda(m^+)^7_7e^{26}.\end{array}\right. \Longrightarrow e^{26}E_6 = qE_6e^{26}.$$

Combing with the above relation $e^{26} = q^{-1}E_6e^{27} - e^{27}E_6$ again, we obtain

$$(E_6)^2E_7 - (q+q^{-1})E_6E_7E_6 + E_7(E_6)^2 = 0.$$

The relations between f_{27} and others F_i, K_i's can be obtained in a similar way. With these relations, we prove that the new quantum group is $U_q(E_7)$. □

3.3. **Inductive construction of $U_q(E_8)$.** We choose the 56-dimensional minimal fundamental representation T_V of $U_q(E_7)$ to construct $U_q(E_8)$. The representation is given by the following Figure 3, which also appeared in [KKM]. This representation also satisfies $E_\beta(f_m) = f_n \iff F_\beta(f_n) = f_m$ and E_i^2 is zero action. Moreover, the data R, R', and the entries we need in $m^\pm$ are given by the following Lemma.

Lemma 3.3. (1) *The minimal polynomial of the braiding PR_{VV} is*

$$(PR_{VV} + q^{-\frac{1}{2}}I)(PR_{VV} + q^{\frac{1}{2}}I)(PR_{VV} - q^{\frac{1}{2}}I)(PR_{VV} - q^{\frac{3}{2}}I) = 0.$$

Thus setting $R = q^{\frac{1}{2}}R_{VV}, R' = -q^{-4}R(PR)^2 + q^{-2}RPR + q^{-2}R$, we have $(PR+I)(PR'-I) = 0$.

(2) *The entries we need in $m^\pm$ are listed as follows*

- *In m^+:*

$$(m^+)_2^1 = -(q-q^{-1})E_7K_1K_2^{\frac{3}{2}}K_3^2K_4^3K_5^{\frac{5}{2}}K_6^2K_7^{\frac{1}{2}},\ (m^+)_2^2 = K_1K_2^{\frac{3}{2}}K_3^2K_4^3K_5^{\frac{5}{2}}K_6^2K_7^{\frac{1}{2}},$$
$$(m^+)_3^2 = -(q-q^{-1})E_6K_1K_2^{\frac{3}{2}}K_3^2K_4^3K_5^{\frac{5}{2}}K_6K_7^{\frac{1}{2}},\ (m^+)_3^3 = K_1K_2^{\frac{3}{2}}K_3^2K_4^3K_5^{\frac{5}{2}}K_6K_7^{\frac{1}{2}},$$
$$(m^+)_4^3 = -(q-q^{-1})E_5K_1K_2^{\frac{3}{2}}K_3^2K_4^3K_5^{\frac{3}{2}}K_6K_7^{\frac{1}{2}},\ (m^+)_4^4 = K_1K_2^{\frac{3}{2}}K_3^2K_4^3K_5^{\frac{3}{2}}K_6K_7^{\frac{1}{2}},$$
$$(m^+)_5^4 = -(q-q^{-1})E_4K_1K_2^{\frac{3}{2}}K_3^2K_4^2K_5^{\frac{3}{2}}K_6K_7^{\frac{1}{2}},\ (m^+)_5^5 = K_1K_2^{\frac{3}{2}}K_3^2K_4^2K_5^{\frac{3}{2}}K_6K_7^{\frac{1}{2}},$$
$$(m^+)_6^5 = -(q-q^{-1})E_3K_1K_2^{\frac{3}{2}}K_3K_4^2K_5^{\frac{3}{2}}K_6K_7^{\frac{1}{2}},\ (m^+)_6^6 = K_1K_2^{\frac{3}{2}}K_3K_4^2K_5^{\frac{3}{2}}K_6K_7^{\frac{1}{2}},$$
$$(m^+)_7^5 = (q^{-1}-q)E_2K_1K_2^{\frac{1}{2}}K_3^2K_4^2K_5^{\frac{3}{2}}K_6K_7^{\frac{1}{2}},\quad (m^+)_7^7 = K_1K_2^{\frac{1}{2}}K_3^2K_4^2K_5^{\frac{3}{2}}K_6K_7^{\frac{1}{2}}, (m^+)_7^6 = 0,$$
$$(m^+)_8^6 = -(q-q^{-1})E_1K_2^{\frac{3}{2}}K_3K_4^2K_5^{\frac{3}{2}}K_6K_7^{\frac{1}{2}},\quad (m^+)_8^8 = K_2^{\frac{3}{2}}K_3K_4^2K_5^{\frac{3}{2}}K_6K_7^{\frac{1}{2}},\ (m^+)_8^7 = 0,$$
$$(m^+)_1^1 = K_1K_2^{\frac{3}{2}}K_3^2K_4^3K_5^{\frac{5}{2}}K_6^2K_7^{\frac{3}{2}},\quad (m^+)_{56}^{56} = K_1^{-1}K_2^{-\frac{3}{2}}K_3^{-2}K_4^{-3}K_5^{-\frac{5}{2}}K_6^{-2}K_7^{-\frac{3}{2}};$$

- *In m^-:*

$$(m^-)_1^2 = \frac{K_1^{-1}K_2^{-\frac{3}{2}}K_3^{-2}K_4^{-3}K_5^{-\frac{5}{2}}K_6^{-2}K_7^{-\frac{1}{2}}F_7}{q^2-1},\quad (m^-)_2^2 = K_1^{-1}K_2^{-\frac{3}{2}}K_3^{-2}K_4^{-3}K_5^{-\frac{5}{2}}K_6^{-2}K_7^{-\frac{1}{2}},$$
$$(m^-)_2^3 = \frac{K_1^{-1}K_2^{-\frac{3}{2}}K_3^{-2}K_4^{-3}K_5^{-\frac{5}{2}}K_6^{-1}K_7^{-\frac{1}{2}}F_6}{q^2-1},\quad (m^-)_3^3 = K_1^{-1}K_2^{-\frac{3}{2}}K_3^{-2}K_4^{-3}K_5^{-\frac{5}{2}}K_6^{-1}K_7^{-\frac{1}{2}},$$
$$(m^-)_3^4 = \frac{K_1^{-1}K_2^{-\frac{3}{2}}K_3^{-2}K_4^{-3}K_5^{-\frac{3}{2}}K_6^{-1}K_7^{-\frac{1}{2}}F_5}{q^2-1},\quad (m^-)_4^4 = K_1^{-1}K_2^{-\frac{3}{2}}K_3^{-2}K_4^{-3}K_5^{-\frac{3}{2}}K_6^{-1}K_7^{-\frac{1}{2}},$$
$$(m^-)_4^5 = \frac{K_1^{-1}K_2^{-\frac{3}{2}}K_3^{-2}K_4^{-2}K_5^{-\frac{3}{2}}K_6^{-1}K_7^{-\frac{1}{2}}F_4}{q^2-1},\quad (m^-)_5^5 = K_1^{-1}K_2^{-\frac{3}{2}}K_3^{-2}K_4^{-2}K_5^{-\frac{3}{2}}K_6^{-1}K_7^{-\frac{1}{2}},$$
$$(m^-)_5^6 = \frac{K_1^{-1}K_2^{-\frac{3}{2}}K_3^{-1}K_4^{-2}K_5^{-\frac{3}{2}}K_6^{-1}K_7^{-\frac{1}{2}}F_3}{q^2-1},\quad (m^-)_6^6 = K_1^{-1}K_2^{-\frac{3}{2}}K_3^{-1}K_4^{-2}K_5^{-\frac{3}{2}}K_6^{-1}K_7^{-\frac{1}{2}},$$
$$(m^-)_6^8 = \frac{K_2^{-\frac{3}{2}}K_3^{-1}K_4^{-2}K_5^{-\frac{3}{2}}K_6^{-1}K_7^{-\frac{1}{2}}F_1}{q^2-1},\quad (m^-)_8^8 = K_2^{-\frac{3}{2}}K_3^{-1}K_4^{-2}K_5^{-\frac{3}{2}}K_6^{-1}K_7^{-\frac{1}{2}}, (m^-)_6^7 = 0,$$
$$(m^-)_8^{10} = \frac{K_2^{-\frac{1}{2}}K_3^{-1}K_4^{-2}K_5^{-\frac{3}{2}}K_6^{-1}K_7^{-\frac{1}{2}}F_2}{q^2-1},\quad (m^-)_{10}^{10} = K_2^{-\frac{1}{2}}K_3^{-1}K_4^{-2}K_5^{-\frac{3}{2}}K_6^{-1}K_7^{-\frac{1}{2}}, (m^-)_8^9 = 0,$$
$$(m^-)_1^1 = K_1^{-1}K_2^{-\frac{3}{2}}K_3^{-2}K_4^{-3}K_5^{-\frac{5}{2}}K_6^{-2}K_7^{-\frac{3}{2}},\ (m^-)_{56}^{56} = K_1K_2^{\frac{3}{2}}K_3^2K_4^3K_5^{\frac{5}{2}}K_6^2K_7^{\frac{3}{2}}.$$

Proof. The proof is similar. The decomposition of this 56-dimensional representation is

$$(0000001) \otimes (0000001) = (0000002) \oplus (0000010) \oplus (1000000) \oplus (0000000),$$

namely, $V \otimes V = V_1 \oplus V_2 \oplus V_3 \oplus V_4$, where $V_i, i = 1, 2, 3, 4$ is the irreducible representation wit highest weight $2\lambda_7, \lambda_6, \lambda_1, 0$ respectively. This means matrix PR_{VV} only has 4 distinct eigenvalues, denoted by $x_i, i = 1, 2, 3, 4$. We set $\triangle_1 = x_1 + x_2 + x_3 + x_4, \triangle_2 = x_1x_2 + x_1x_3 + x_2x_3 + (x_1 + x_2 + x_3)x_4, \triangle_3 = x_1x_2x_3 + x_1x_2x_4 + x_1x_3x_4 + x_2x_3x_4, \triangle_4 = x_1x_2x_3x_4$, and $\mathcal{N} = (PR_{VV} - x_1I)(PR_{VV} - x_2I)(PR_{VV} - x_3I)\,(PR_{VV} - x_4I)$.

We find nonzero entries at rows (12) and (21) in matrix $PR_{VV} - x_iI$ are $(PR_{VV} - x_iI)^{12}_{12} = -x_i, (PR_{VV} - x_iI)^{12}_{21} = q^{\frac{1}{2}}, (PR_{VV} - x_iI)^{21}_{12} = q^{\frac{1}{2}}, (PR_{VV} - x_iI)^{21}_{21} = q^{\frac{1}{2}}(q - q^{-1}) - x_i$. So we obtain the entries at row $(1, 2)$ and columns $(1, 2)$ and $(2, 1)$ in matrix $\mathcal{N}$ are

$$\mathcal{N}^{12}_{12} = \Delta_4 + q\Delta_2 - q^{\frac{3}{2}}(q - q^{-1})\Delta_1 + q^2 + q^2(q - q^{-1})^2,$$
$$\mathcal{N}^{12}_{21} = -\Delta_3 + q^{\frac{1}{2}}(q - q^{-1})\Delta_2 - [q + q(q - q^{-1})^2]\Delta_1 + q^{\frac{3}{2}}(q - q^{-1})^3 + 2q^{\frac{3}{2}}(q - q^{-1}).$$

We will consider the entries at rows $(5, 11)$ and $(11, 5)$. From the Figure 3, we find that the basis f_5 can be raised by the roots α_2,α_3 and $\alpha_1 + \alpha_3$, but the basis f_{11} is lowered by the roots α_4,$\alpha_3 + \alpha_4$ and $\alpha_2 + \alpha_4$. This fact means that raising f_5 and lowering f_{11} can't be done by the same roots, so the only nonzero entry at row $(5, 11)$ in matrix R_{VV} is $R_{VV}{}^{5,11}_{5,11} = q^{\frac{1}{2}}$. We also obtain that the only nonzero entry at row $(11, 5)$ in matrix R_{VV} is $R_{VV}{}^{11,5}_{11,5} = q^{\frac{1}{2}}$ by the same analysis. Then the nonzero entries at rows $(5, 11)$ and $(11, 5)$ in matrix $PR_{VV} - x_iI$ are $(PR_{VV} - x_iI)^{5,11}_{5,11} = -x_i, (PR_{VV} - x_iI)^{5,11}_{11,5} = q^{\frac{1}{2}}, (PR_{VV} - x_iI)^{11,5}_{5,11} = q^{\frac{1}{2}}, (PR_{VV} - x_iI)^{11,5}_{11,5} = -x_i$. Then the entries at row $(5, 11)$ and columns $(5, 11)$ and $(11, 5)$ in matrix $\mathcal{N}$ are $\mathcal{N}^{5,11}_{5,11} = \Delta_4 + q\Delta_2 + q^2, \mathcal{N}^{5,11}_{11,5} = -q^{\frac{1}{2}}\Delta_3 - q^{\frac{3}{2}}\Delta_1$.

These entries yield the following equations

$$\begin{cases} \Delta_4 + q\Delta_2 - q^{\frac{3}{2}}(q - q^{-1})\Delta_1 + q^2 + q^2(q - q^{-1})^2 = 0, \\ -\Delta_3 + q^{\frac{1}{2}}(q - q^{-1})\Delta_2 - [q + q(q - q^{-1})^2]\Delta_1 + q^{\frac{3}{2}}(q - q^{-1})^3 + 2q^{\frac{3}{2}}(q - q^{-1}) = 0, \\ \Delta_4 + q\Delta_2 + q^2 = 0, \\ -q^{\frac{1}{2}}\Delta_3 - q^{\frac{3}{2}}\Delta_1 = 0. \end{cases}$$

Solving it, we get these eigenvalues are $-q^{-\frac{1}{2}}, \pm q^{\frac{1}{2}}, q^{\frac{3}{2}}$. Then the minimal polynomial of PR_{VV} is $(PR_{VV} + q^{-\frac{1}{2}}I)(PR_{VV} + q^{\frac{1}{2}}I)(PR_{VV} - q^{\frac{1}{2}}I)(PR_{VV} - q^{\frac{3}{2}}I) = 0$. □

With these, we have the following

Theorem 3.3. *With the quantum normalization constant $\lambda = q^{-\frac{1}{2}}$, identify e^{56}, f_{56} and $(m^+)^{56}_{56}c^{-1}$ with the additional simple root vectors E_8, F_8, and the group-element K_8, then the new quantum group $U(V^\vee(R', R_{21}^{-1}), U_q^{\widetilde{ext}}(E_7), V(R', R)) = U_q(E_8)$ with $K_i^{\frac{1}{2}}$ adjoined.*

Proof. The cross relation can be obtained easily in view of the relations in Theorem 2.2, so we only focus on the q-Serre relations between e^{56} and E_i. E_1, E_2 appear in $(m^+)^6_8$, $(m^+)^5_7$, respectively, then according to Theorem 2.2, we obtain

$$\left.\begin{array}{l}(m^+)_8^6 = -(q-q^{-1})E_1(m^+)_8^8,\\ e^{56}(m^+)_8^6 = \lambda R_{656}^{656}(m^+)_8^6 e^{56} = \lambda(m^+)_8^6 e^{56},\\ e^{56}(m^+)_8^8 = \lambda R_{856}^{856}(m^+)_8^6 e^{56} = \lambda(m^+)_8^8 e^{56},\end{array}\right\} \Longrightarrow E_8E_1 = E_1E_8.$$

$$\left.\begin{array}{l}(m^+)_7^5 = -(q-q^{-1})E_2(m^+)_7^7,\\ e^{56}(m^+)_7^5 = \lambda R_{556}^{556}(m^+)_7^5 e^{56} = \lambda(m^+)_7^5 e^{56},\\ e^{56}(m^+)_7^7 = \lambda R_{756}^{756}(m^+)_7^7 e^{56} = \lambda(m^+)_7^7 e^{56},\end{array}\right\} \Longrightarrow E_8E_2 = E_2E_8.$$

We observed that other E_i's locate in the minor entries $(m^+)_{i+1}^i$ from the above Lemma, then the following relations can be deduced from Theorem 2.2.

$$\left.\begin{array}{l}e^{56}(m^+)_2^1 = \lambda R_{156}^{156}(m^+)_2^1 e^{56} + \lambda R_{255}^{156}(m^+)_2^2 e^{55},\\ e^{56}(m^+)_{i+1}^i = \lambda R_{i56}^{i56}(m^+)_{i+1}^i e^{56}, 2 \le i \le 5,\\ (m^+)_{i+1}^i = -(q-q^{-1})E_j(m^+)_{i+1}^{i+1}, i+j = 8,\\ e^{56}(m^+)_k^k = \lambda R_{k56}^{k56}(m^+)_k^k e^{56}, \forall k,\\ R_{255}^{255} = q^{-1}, R_{2,55}^{1,56} = q^{-1}(q-q^{-1}),\\ R_{156}^{156} = q^{-1}, R_{i56}^{i56} = 1, 2 \le i \le 8,\end{array}\right\} \Longrightarrow \left\{\begin{array}{l}E_8E_j = E_jE_8,\ 3 \le j \le 6,\\ e^{55} = qE_7e^{56} - e^{56}E_7.\end{array}\right.$$

According to $R'^{56,55}_{56,55} = q^{-1} - q^{-3}$, and $R'^{56,55}_{55,56} = q^{-2}$, then $e^{55}e^{56} = R'^{56\,55}_{a\ b}e^a e^b = (q^{-1}-q^{-3})e^{56}e^{55}+q^{-2}e^{55}e^{56} \Longrightarrow e^{55}e^{56} = q^{-1}e^{56}e^{55}$. Combining with $e^{55} = qE_7e^{56}-e^{56}E_7$, we obtain

$$(E_8)^2E_7 - (q+q^{-1})E_8E_7E_8 + E_7(E_8)^2 = 0.$$

On the other hand, we can get the relations between e^{55} and E_7 by the following relations.

$$\left.\begin{array}{l}m^{+1}_{\ 2} = -(q-q^{-1})E_7m^{+2}_{\ 2},\\ e^{55}m^{+1}_{\ 2} = \lambda R_{1,55}^{1,55}m^{+1}_{\ 2}e^{55},\\ e^{55}m^{+2}_{\ 2} = \lambda R_{2,55}^{2,55}m^{+2}_{\ 2}e^{55},\end{array}\right\} \Longrightarrow e^{55}E_7 = q^{-1}E_7e^{55}.$$

Combining with $e^{55} = qE_7e^{56} - e^{56}E_7$ again, we obtain

$$(E_7)^2E_8 - (q+q^{-1})E_7E_8E_7 + E_8(E_7)^2 = 0.$$

With these relations, we prove that the new quantum group is $U_q(E_8)$.

This completes the proof. □

Figure 2. $U_q(E_6)$'s 27-dimensional minimal fundamental representation



Figure 3. $U_q(E_7)$'s 56-dimensional minimal fundamental representation



4. Majid's double-bosonization and Rosso's quantum shuffle

Up to now, we have obtained the inductive constructions of $U_q(\mathfrak{g})$'s for all finite-dimensional complex simple Lie algebras $\mathfrak{g}$, which can be expressed in a unified form by the following

Proposition 4.1. *Let T_V be a p-dimensional irreducible representation of $U_q(\mathfrak{g})$ with highest weight $-\mu$, $(a_{ij})_{n\times n}$ the Cartan matrix of $\mathfrak{g}$, ν the weight of central element c^{-1}. Then the resulted quantum group $U(V^\vee(R', R_{21}^{-1}), \widetilde{U_q^{ext}(\mathfrak{g})}, V(R', R))$ is of higher-one rank, whose Cartan matrix is obtained from $(a_{ij})_{n\times n}$ by adding a row and a column with: $a_{i,n+1} = \frac{2(\alpha_i,\mu)}{(\alpha_i,\alpha_i)}$, $a_{n+1,i} = \frac{2(\mu,\alpha_i)}{(\mu,\mu)+(\nu,\nu)}$, and ν is orthogonal to μ and $\alpha_i, i = 1, \cdots, n$.*

Proof. Combining with the inductive constructions obtained in [HH1, HH2] and the results in section 3 of this paper, we know that the new additional group-like element K_{n+1} is $(m^+)_p^p c^{-1}$. From the expression of $(m^+)_p^p$ in each case, we observe that the corresponding weight of $(m^+)_p^p$ is μ, namely $(m^+)_p^p = K_\mu$, so $K_{n+1} = K_{\mu+\nu}$. Every simple root vector E_j usually locates in the minor entries $(m^+)_{i+1}^i$, where $(m^+)_{i+1}^i = a_i E_j (m^+)_{i+1}^{i+1}$, $a_i \in k[q, q^{-1}]$. According to the cross relations in Theorem 2.2, we obtain the following relations between e^i and e^{i-1}.

$$\begin{cases} e^i(m^+)_{i+1}^i = \lambda R_{ab}^{i\,i}(m^+)_{i+1}^a e^b = \lambda R_{ii}^{ii}(m^+)_{i+1}^i e^i + \lambda R_{i+1\,i-1}^{i\;\;i}(m^+)_{i+1}^{i+1} e^{i-1}, \\ e^i(m^+)_{i+1}^{i+1} = \lambda R_{i+1\,i}^{i+1\,i}(m^+)_{i+1}^{i+1} e^i, \\ e^{i-1}(m^+)_{i+1}^{i+1} = \lambda R_{i+1\,i-1}^{i+1\,i-1}(m^+)_{i+1}^{i+1} e^{i-1}. \end{cases}$$

Combining with $(m^+)_{i+1}^i = a_i E_j (m^+)_{i+1}^{i+1}$, we get $e^{i-1} = a_i \frac{R_{i+1\,i-1}^{i+1\,i-1}}{R_{i+1\,i-1}^{i\;\;i}}(e^i E_j - \frac{R_{ii}^{ii}}{R_{i+1\,i}^{i+1\,i}} E_j e^i)$. Then $\mathrm{weight}(e^{i-1}) = \mathrm{weight}(e^i) + \alpha_j$, combining with $e^k \cdot c^{-1} = \lambda^{-1} e^k$ for any k, so $(\nu, \alpha_j) = 0$. On the other hand, we know that the explicit form of the additional $K_{\mu+\nu}$ in each inductive construction, so we can obtain the specific form of ν, and observe that ν is orthogonal to μ. □

With these explicit inductive constructions, we know that both double bosonization [M1] and quantum shuffle in [Ro] have the same application for the inductive constructions of $U_q(\mathfrak{g})$'s for the finite-dimensional complex simple Lie algebras. We will give the specific form of weight μ and ν, and the Dynkin diagram in each case, which makes the readers understand intuitively. The imaginary line and the filled circle mean that the original Dynkin diagram extends to the new added simple root with an arrow pointing to the shorter of the two roots in each extended Dynkin diagram.

(1) The A_n series.

Take $\mathfrak{g} = \mathfrak{sl}_n$ and V the fundamental representation with $\mu = -\lambda_{n-1}$, where $\lambda_{n-1} = \sum_{i=1}^{n-1} \frac{i}{n}\alpha_i = \frac{1}{n}\sum_{i=1}^{n-1} \varepsilon_i - \frac{n-1}{n}\varepsilon_n$. $\nu = \frac{1}{n}\sum_{i=1}^{n} \varepsilon_i - \varepsilon_{n+1}$, then we get $\mathfrak{sl}_{n+1}$ ([HH1]).


α_1 α_2 α_{n-2} α_{n-1}
$\varepsilon_1 - \varepsilon_2$ $\varepsilon_2 - \varepsilon_3$ $\varepsilon_{n-2} - \varepsilon_{n-1}$ $\varepsilon_{n-1} - \varepsilon_n$ $\varepsilon_n - \varepsilon_{n+1}$

(2) The B_n series.

B_2 can be constructed from A_1 in [HH1]. Then starting from $\mathfrak{g} = \mathfrak{so}_{2n+1}$ and V is the fundamental representation with $\mu = -\lambda_n$, where $\lambda_n = \sum_{i=1}^{n} \alpha_i = \varepsilon_n$. $\nu = \varepsilon_{n+1}$, then we get $\mathfrak{so}_{2n+3}$ ([HH1]).


α_1 α_2 α_{n-1} α_n
ε_1 $\varepsilon_2 - \varepsilon_1$ $\varepsilon_{n-1} - \varepsilon_{n-2}$ $\varepsilon_n - \varepsilon_{n-1}$ $\varepsilon_{n+1} - \varepsilon_n$

(3) The C_n series.

C_3 can be constructed from A_2 in [HH1]. Then starting from $\mathfrak{g} = \mathfrak{sp}_{2n}$ and V is the fundamental representation with $\mu = -\lambda_n$, where $\lambda_n = \frac{1}{2} + \sum_{i=2}^{n} \alpha_i = \varepsilon_n$. $\nu = \varepsilon_{n+1}$, then we get $\mathfrak{sp}_{2n+2}$ ([HH1]).


α_1 α_2 α_{n-1} α_n
$2\varepsilon_1$ $\varepsilon_2 - \varepsilon_1$ $\varepsilon_{n-1} - \varepsilon_{n-2}$ $\varepsilon_n - \varepsilon_{n-1}$ $\varepsilon_{n+1} - \varepsilon_n$

(4) The D_n series.

D_4 can be constructed from A_3 in [HH1]. Then starting from $\mathfrak{g} = \mathfrak{so}_{2n}$ and V is the fundamental representation with $\mu = -\lambda_n$, where $\lambda_n = \frac{1}{2}(\alpha_1 + \alpha_2) + \sum_{i=3}^{n} \alpha_i = \varepsilon_n$. $\nu = \varepsilon_{n+1}$, then we get $\mathfrak{so}_{2n+2}$ ([HH1]).


$\alpha_1 = \varepsilon_2 - \varepsilon_1$
α_3 α_{n-1} α_n
$\varepsilon_3 - \varepsilon_2$ $\varepsilon_{n-1} - \varepsilon_{n-2}$ $\varepsilon_n - \varepsilon_{n-1}$ $\varepsilon_{n+1} - \varepsilon_n$
$\alpha_2 = \varepsilon_2 + \varepsilon_1$

(5) The case of E_6.

Take $\mathfrak{g} = \mathfrak{so}_{10}$ and V the 16-dimensional half-spin representation with $\mu = -\lambda_5$, where $\lambda_5 = \frac{1}{2}\alpha_1 + \alpha_2 + \frac{3}{2}\alpha_3 + \frac{3}{4}\alpha_4 + \frac{5}{4}\alpha_5 = \frac{1}{2}\sum_{i=1}^{5} \varepsilon_i$. $\nu = \frac{1}{2}(\varepsilon_7 + \varepsilon_8 - \varepsilon_6)$, then we get E_6.


$\alpha_4 = \varepsilon_4 - \varepsilon_5$
α_1 α_2 α_3 α_5
$\varepsilon_1 - \varepsilon_2$ $\varepsilon_2 - \varepsilon_3$ $\varepsilon_3 - \varepsilon_4$ $\varepsilon_4 + \varepsilon_5$ $\frac{1}{2}(\varepsilon_7 + \varepsilon_8 - (\varepsilon_1 + \cdots + \varepsilon_6))$

(6) The case of E_7.

Take $\mathfrak{g} = E_6$ and V the 27-dimensional fundamental representation with $\mu = -\lambda_6$, $\lambda_6 = \frac{2}{3}\alpha_1 + \alpha_2 + \frac{4}{3}\alpha_3 + 2\alpha_4 + \frac{5}{3}\alpha_5 + \frac{4}{3}\alpha_6 = \frac{1}{3}\varepsilon_8 - \frac{1}{3}(\varepsilon_6 + \varepsilon_7) + \varepsilon_5$, and $\nu = \frac{1}{3}\varepsilon_8 + \frac{2}{3}\varepsilon_6 - \frac{1}{3}\varepsilon_7$, then we get E_7.


$\alpha_2 = \varepsilon_2 + \varepsilon_1$
α_6 α_5 α_4 α_3 α_1
$\varepsilon_6 - \varepsilon_5$ $\varepsilon_5 - \varepsilon_4$ $\varepsilon_4 - \varepsilon_3$ $\varepsilon_3 - \varepsilon_2$ $\varepsilon_2 - \varepsilon_1$ $\frac{1}{2}(\varepsilon_1 + \varepsilon_8 - (\varepsilon_2 + \cdots + \varepsilon_7))$

(7) The case of E_8.

Take $\mathfrak{g} = E_7$ and V the 56-dimensional fundamental representation with $\mu = -\lambda_7$, $\lambda_7 = \alpha_1 + \frac{3}{2}\alpha_2 + 2\alpha_3 + 3\alpha_4 + \frac{5}{2}\alpha_5 + 2\alpha_6 + \frac{3}{2}\alpha_7 = \frac{1}{2}\varepsilon_8 - \frac{1}{2}\varepsilon_7 + \varepsilon_6$, and $\nu = \frac{1}{2}(\varepsilon_8 + \varepsilon_7)$, then we get E_8.


$\alpha_2 = \varepsilon_2 + \varepsilon_1$
α_7 α_6 α_5 α_4 α_3 α_1
$\varepsilon_7 - \varepsilon_6$ $\varepsilon_6 - \varepsilon_5$ $\varepsilon_5 - \varepsilon_4$ $\varepsilon_4 - \varepsilon_3$ $\varepsilon_3 - \varepsilon_2$ $\varepsilon_2 - \varepsilon_1$ $\frac{1}{2}(\varepsilon_1 + \varepsilon_8 - (\varepsilon_2 + \cdots + \varepsilon_7))$

(8) The case of F_4.

Take $\mathfrak{g} = \mathfrak{so}_7$ and V the 8-dimensional spin representation with $\mu = -\lambda_3$, $\lambda_3 = \frac{1}{2}\alpha_1 + \alpha_2 + \frac{3}{2}\alpha_3 = \frac{1}{2}\varepsilon_1 + \frac{1}{2}\varepsilon_2 + \frac{1}{2}\varepsilon_3$, and $\nu = \frac{1}{2}\varepsilon_4$, then we get F_4 (see [HH2]).


α_1 α_2 α_3
$\varepsilon_3 - \varepsilon_2$ $\varepsilon_2 - \varepsilon_1$ ε_1 $\frac{1}{2}(\varepsilon_4 - \varepsilon_1 - \varepsilon_2 - \varepsilon_3)$

(9) The case of G_2.

Take $\mathfrak{g} = \mathfrak{sl}_2$ and V the 4-dimensional spin $\frac{3}{2}$ representation with $\mu = -3\lambda_1$, $\lambda_1 = \frac{1}{2}\alpha_1 = \frac{1}{2}(\varepsilon_1 - \varepsilon_2)$, and $\nu = -\frac{1}{2}(\varepsilon_1 + \varepsilon_2) + \varepsilon_3$, then we get G_2 (see [HH2]).



REFERENCES

[AS1] N. Andruskiewitsch and H.-J. Schneider, On the classification of finite-dimensional pointed Hopf algebras, *Ann. of Math. (2) 171* (1) (2010), 375–417.

[AS2] N. Andruskiewitsch and H.-J. Schneider, Pointed Hopf algebras, New directions in Hopf algebras, 1–68, Math. Sci. Res. Inst. Publ., 43, Cambridge Univ. Press, Cambridge, 2002.

[Br] T. Bridgeland, Quantum groups via Hall algebras of complexes, *Ann. of Math. (2) 177 (2)* (2013), 739–759.

[CL] M. Cuntz and S. Lentner, A simplicial complex of Nichols algebras, arXiv:1503.08117.

[D] V.G. Drinfeld, Hopf algebras and the quantum Yang-Baxter equation, *Dokl. Akad. Nauk SSSR 283 (5)* (1985), 1060–1064.

[FRT] L.D. Faddeev, N.Yu. Reshetikhin and L.A. Takhtajan, Quantization of Lie groups and Lie algebras, *Leningrad Math. J. 1* (1990), 193–225.

[FR] X. Fang and M. Rosso, Multi-brace cotensor Hopf algebras and quantum groups, arXiv:1210.3096v1.

[G] J.E. Grabowski, Braided enveloping algebras associated to quantum parabolic subalgebras, *Comm. Algebra 39* (10) (2011), 3491–3514.

[HS] I. Heckenberger and H.-J. Schneider, Yetter-Drinfeld modules over bosonizations of dually paired Hopf algebras, *Adv. Math. 244* (2013), 354–394.

[HH1] Hongmei Hu and Naihong Hu, *Double-bosonization and Majid's conjecture, (I): rank-inductions of ABCD*, J. Math. Phys. **56** (11) (2015), 111702-1, 16 pp.

[HH2] —, *Double-bosonization and Majid's conjecture, (II): cases of irregular R-matrices and type-crossings of F_4, G_2*, arXiv:1512.08712.

[HH3] —, *Double-bosonization and Majid's conjecture, (IV): type-crossings from A to BCD*, Sci. China Math. (2016) (to appear), arXiv: 1505.02620.

[HLR] N.H. Hu, Y.N. Li and M. Rosso, Multi-parameter quantum groups via quantum quasi-symmetric algebras, arXiv:1307.1381.

[Ji] M. Jimbo, A q-difference analog of $U(g)$ and the Yang-Baxter equation, *Lett. Math. Phys. 10 (1)* (1985), 63–69.

[K] A. Klimyk and K. Schmdgen, Quantum Groups and their Representations, *Springer-Verlag, Berlin Heidelberg*, 1997.

[KKM] J.D. Kim, I.G. Koh and Zhong-Qi Ma, Quantum R-matrix for E_7 and F_4 groups, *J. Math. Phys. 32* (1991), 845–856.

[KM] I.G. Koh, Zhong-Qi Ma, Exceptional quantum groups, *Phys. Lett. B 234 (4)* (1990), 480–486.

[L1] G. Lusztig, Canonical bases arising from quantized enveloping algrbras, *J. Amer. Math. Soc. 3* (1990), 447–498.

[L2] G. Lusztig, Introduction to Quantum Groups, *Progress in Math. 110*, Birkhäuser, Boston (1993).

[M1] S. Majid, Double-Bosonization of braided groups and the construction of $U_q(\mathfrak{g})$, *Math. Proc. Cambridge Philos. Soc. 125* (1999), 151–192.

[M2] S. Majid, Braided groups, *J. Pure and Applied Algebra 86* (1993), 187–221.

[M3] S. Majid. Algebras and Hopf algebras in braided categories, *Lecture Notes in Pure and Appl. Math. 158 (1994)*, 55–105.

[M4] S. Majid, Braided momentum in the q-Poincare group, *J. Math. Phys 34* (1993), 2045–2058.

[M5] S. Majid, More examples of bicrossproduct and double cross product Hopf algebras, *Isr. J. Math 72* (1990), 133–148.
[M6] S. Majid, New quantum groups by double-bosonization, *Czechoslovak J. Phys. 47 (1)* (1997), 79–90.
[M7] S. Majid, Braided geometry and inductive construction of Lie algebras and quantum groups, arXiv:q-alg/9611006v1 (1996).
[Ra] D. Radford, Hopf algebras with projection, *J. Algebra 92* (1985), 322–347.
[Ri] C. Ringel, Hall algebras and quantum groups, *Invent. Math. 101* (1990), 583–592.
[Ro] M. Rosso, Quantum groups and quantum shuffles, *Invent. Math. 133 (2)* (1998), 399–416.
[So] Y. Sommerhäuser, *Deformed enveloping algebras*, New York J. Math. **2** (1996), 35–58.

Department of Mathematics, Shanghai Key Laboratory of Pure Mathematics and Mathematical Practice, East China Normal University, Minhang Campus, Dong Chuan Road 500, Shanghai 200241, PR China

E-mail address: hmhu0124@126.com

Department of Mathematics, Shanghai Key Laboratory of Pure Mathematics and Mathematical Practice, East China Normal University, Minhang Campus, Dong Chuan Road 500, Shanghai 200241, PR China

E-mail address: nhhu@math.ecnu.edu.cn